

# PRODUCT INFORMATION



PERFORMANCE MINERALS™

## ZINPRO 100


### ZINC METHIONINE COMPLEX

A nutritional feed ingredient for livestock and poultry. When used as a commercial feed ingredient it must be declared as zinc methionine complex. Association of American Feed Control Officials (AAFCO) No. 57.151. International Feed Number (IFN) 6-16-293. Zinc Methionine Complex. U.S. Patent No. 4,764,633 and 5,430,164. Canada Reg. No. 990582.

#### TYPICAL ANALYSIS

Zinc .....	10.0%
Methionine .....	20.0%
Protein .....	11.8%
Fat .....	0.0%
Fiber .....	15.0%
Ash .....	20.0%
Salt .....	1.2%

#### MOLECULAR STRUCTURE



#### PHYSICAL DESCRIPTION

A light brown granular powder with characteristic methionine-like odor. ZINPRO® 100 weighs approximately 35 lb/cu. ft. and is packaged in 50 lb multiwall bags.

#### FEEDING INSTRUCTIONS

**SWINE\*:** Add 0.8 lb (363 grams) per ton of complete diet.

**LAYING HENS, BROILERS AND TURKEYS:** Add 0.8 lb (363 grams) per ton of complete diet.

**DAIRY CATTLE:** Feed 3.6 grams per head daily, or 1 lb per 125 head daily.

**BEEF CATTLE:** Feed 3.6 grams per head daily, or 1 lb per 125 head daily.

**HORSES:** Feed 3.6 grams per head daily.

**SHEEP\*:** Feed 0.4 grams per head daily, or 1 lb per 1,250 head daily.

**GOATS\*:** Feed 0.5 grams per head daily, or 1 lb per 900 head daily.

\* Double recommended levels during periods of reduced feed intake, or high production stress.

#### TOXICITY AND SAFE HANDLING

When correctly used, there is no toxicity hazard in the use of ZINPRO 100. Store in a clean and dry environment. Follow label directions and rotate inventory to ensure fresh product.